

LA-UR-01:3664

Approved for public release;

distribution is unlimited

Title: Towards a Calculus of Biological
Networks

Author(s): Henning S. Mortveit, D-2
Christian M. Reidys, D-2

Journal: Z. Phys. Chem.
Oldenbourg Wissenschaftsverlag
München
Germany

Journal data: 216 (2002) 235–247

Status: Printed

LOS ALAMOS

NATIONAL LABORATORY

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. The Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

TOWARDS A CALCULUS OF BIOLOGICAL NETWORKS

H.S. MORTVEIT AND C.M. REIDYS

ABSTRACT. In this paper we present a new framework for studying the dynamics of biological networks. A specific class of dynamical systems, *Sequential Dynamical Systems* (SDS), is introduced. These systems allow one to investigate the interplay between structural properties of the network and its phase space. We will show in detail how to find a reduced system that captures key features of a given system. This reduction is based on a special graph-theoretic relation between the two networks. We will study the reduction of SDS over n -cubes in detail and we will present several examples.

1. INTRODUCTION

Biological networks, like metabolic or regulatory networks, as well as networks in general, can be considered as undirected or directed graphs in which the vertices have states that depend on the states of their corresponding adjacencies. One typically has some information about how state transitions of the respective vertices occur, but it is extremely difficult to analyze the global dynamics of the networks as interactions among the vertices occur. Additionally, it appears to be a generic feature of biological networks that these interactions take place sequentially.

Accordingly, a mathematical framework designed for the analysis of the dynamics of biological networks should explicitly take into account: scheduling, the properties of the vertices as state transition functions, and the interconnection scheme, i.e., the graph itself.

Date: June, 2001.

Key words and phrases. sequential dynamical systems, graph morphisms, phase space embedding.

In the following we will introduce a new class of dynamical systems called sequential dynamical systems, or SDS for short [5, 10]. SDS are basically comprised of *i*) state transition rules, *ii*) an undirected graph where each vertex has a state, and *iii*) a permutation of the vertices in the graph giving the update order. Thus an SDS is a dynamical system of the form $\phi : \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$, where $\mathbb{F}_2 = \{0, 1\}$ is the field with two elements.

The question of obtaining information about a network thus becomes a question of understanding the phase space structure of the SDS ϕ . For some results on, e.g., reversibility/invertibility and fixed points we refer to [2, 3, 5, 10, 11].

The phase space of an SDS will usually consist of more than one attractor. Thus a time series will only visit certain parts of phase space. Likewise for networks: there will typically be valid states or regimes that are never realized under the time evolution. It seems fair to believe that constructing a “reduced” network that produces the same dynamics as the original network in the essential regimes and “throws away” the non-essential regimes should allow for more insight and has an obvious computational advantage.

In the following we will show how to reduce the phase space of certain SDS ϕ over a graph Y . We will do so by considering an SDS ψ over a smaller graph Z for which there is a covering map (locally bijective graph morphism) $p : Y \rightarrow Z$. In this construction the state transition functions of corresponding Y and Z vertices are identical, and the dynamics of the SDS ψ represents the essential parts of the dynamics of ϕ . In fact, there can be several covering maps $p_i : Y \rightarrow Z_i$, each of which gives rise to a reduced system. This may be viewed as factorization of the SDS over Y into factors which are SDS over Z_i .

Let Y be a labeled graph with vertex-set $v[Y] = \mathbb{N}_n = \{1, 2, 3, \dots, n\}$, which we write as $Y < K_n$, where K_n is the complete graph on n vertices. The edge-set of Y is denoted by $e[Y]$. Let $S_{1,Y}(i)$ be the set of Y -vertices that are adjacent to vertex i , let $\delta_i = |S_{1,Y}(i)|$ and let $d = \max_{i \in \mathbb{N}_n} \delta_i$. The increasing sequence of elements of $S_{1,Y}(i)$

preceded by i is denoted by

$$(1.1) \quad \tilde{B}_{1,Y}(i) = (i, j_1, \dots, j_{\delta_i}).$$

To each vertex i we associate a state $x_i \in \mathbb{F}_2$, and we write $x = (x_1, x_2, \dots, x_n)$ for the system state. For each $k = 1, \dots, d+1$ we have a symmetric function $f : \mathbb{F}_2^k \rightarrow \mathbb{F}_2$, and for each vertex i we introduce a map

$$(1.2) \quad \text{proj}_Y[i] : \mathbb{F}_2^n \rightarrow \mathbb{F}_2^{\delta_i+1}, \quad (x_1, \dots, x_n) \mapsto (x_i, x_{j_1}, \dots, x_{j_{\delta_i}}).$$

The map projects from the full n -tuple x down to the states vertex i needs for updating its state. For each $i \in \mathbb{N}_n$ there is a Y -local map $F_{i,Y} : \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ given by

$$(1.3) \quad \begin{aligned} y_i &= f_{\delta_i+1} \circ \text{proj}_Y[i], \\ F_{i,Y}(x) &= (x_1, \dots, x_{i-1}, y_i(x), x_{i+1}, \dots, x_n). \end{aligned}$$


The function $F_{i,Y}$ updates the state of vertex i and leaves all other states fixed. We refer to the sequence $(F_{i,Y})_i$ as F_Y . Note that for each graph $Y < K_n$ a sequence $(f_k)_{1 \leq k \leq n}$ induces a sequence F_Y , i.e., we have a map $\{Y < K_n\} \rightarrow \{F_Y\}$. We define the map $[F_Y, \cdot] : S_n \rightarrow \text{Map}(\mathbb{F}_2^n, \mathbb{F}_2^n)$ by

$$(1.4) \quad [F_Y, \pi] = \prod_{i=1}^n F_{\pi(i),Y},$$

where product denotes ordinary function composition.

Definition 1 (Sequential Dynamical System). Let $Y < K_n$, let $(f_k)_k$ with $1 \leq k \leq d(Y) + 1$ be a sequence of symmetric functions, and let $\pi \in S_n$. The sequential dynamical system (SDS) over Y induced by $(f_k)_k$ with respect to the ordering π is $[F_Y, \pi]$.

We call an SDS homogeneous if it is induced by a sequence of local symmetric functions of the form $(f_k)_k = (B_k)_k$ where B is a Boolean function like, e.g., parity which returns the sum of its arguments modulo 2.

FIGURE 1. The circle graph on 4 vertices, Circ_4 .

Example 1. Let $Y = \text{Circ}_4$ as shown in figure 1. With the parity function, i.e., $\text{par}_3 : \mathbb{F}_2^3 \rightarrow \mathbb{F}_2$ defined by $\text{par}_3(x_1, x_2, x_3) = \sum_i x_i \pmod 2$, update order $(1, 2, 3, 4)$ and initial state $(1, 1, 0, 0)$ we get

$$\text{Par}_1(1, 1, 0, 0) = (0, 1, 0, 0),$$

$$\text{Par}_2 \circ \text{Par}_1(1, 1, 0, 0) = (0, 1, 0, 0),$$

$$\text{Par}_3 \circ \text{Par}_2 \circ \text{Par}_1(1, 1, 0, 0) = (0, 1, 1, 0),$$

$$\text{Par}_4 \circ \text{Par}_3 \circ \text{Par}_2 \circ \text{Par}_1(1, 1, 0, 0) = (0, 1, 1, 1).$$

and thus $[\mathbf{Par}_{\text{Circ}_4}, (1, 2, 3, 4)](1, 1, 0, 0) = (0, 1, 1, 1)$.

Since phase space for an SDS is finite we may identify it with a finite unicyclic digraph.

Definition 2. The digraph $\Gamma[F_Y, \pi]$ associated to the SDS $[F_Y, \pi]$ is the directed graph having vertex-set \mathbb{F}_2^n and directed edges $\{(x, [F_Y, \pi](x)) \mid x \in \mathbb{F}_2^n\}$.

A subgroup $H < Q_2^n$ will under certain conditions induce a locally bijective map $\phi : Q_2^n \rightarrow H \setminus Q_2^n$. The graph $H \setminus Q_2^n$ has vertex set Q_2^n/H (factor space) and two vertices \bar{u}, \bar{v} are adjacent iff there are elements $u \in \bar{u}$ and $v \in \bar{v}$ such that $\{u, v\} \in e[Q_2^n]$.

This paper is organized as follows. First we introduce the the concept of graph coverings and locally bijective graph morphisms. Assuming we have a graph covering $\phi : Y \rightarrow Z$, we show how the phase space of an SDS over Z is embedded into the phase space of an SDS over Y where the same local functions are used. Finally, we develop a criterion based on local bijectivity and subgroup structure of the n -cube Q_2^n for the existence of covering maps of the form $\phi : Q_2^n \rightarrow Z = H \setminus Q_2^n$, $H < Q_2^n$.

2. FACTORIZATION

Recall that a morphism between graphs Y and Y' is a pair $\phi = (\phi_1, \phi_2)$ with $\phi_1 : v[Y] \rightarrow v[Y']$ and $\phi : e[Y] \rightarrow e[Y']$ such that

$$\forall e = \{i, j\} \in e[Y] : \quad \phi_2(e) = \{\phi_1(i), \phi_1(j)\}.$$

Thus, adjacent vertices in Y are mapped to *i*) adjacent vertices in Y' or *ii*) to the same vertex in Y' . A morphism of directed graphs also preserves the direction of edges.

A graph morphism $\phi : Y \rightarrow Y'$ is *locally bijective (surjective)* if

$$\forall i \in v[Y] : \phi|_{B_{1,Y}(i)} : B_{1,Y}(i) \rightarrow B_{1,Y'}(\phi(i))$$

is bijective (surjective). Note that a locally bijective graph morphism does not have to be bijective as the following example shows.

Example 2. As an example of a locally bijective graph morphism we have $\phi : Q_2^3 \rightarrow K_4$, see figure 2. The map ϕ_1 is defined by $\phi_1(\{0, 7\}) =$





FIGURE 2. The graphs Q_2^3 and K_4 .

$\{1\}$, $\phi_1(\{1, 6\}) = \{2\}$, $\phi_1(\{2, 5\}) = \{3\}$, $\phi_1(\{3, 4\}) = \{4\}$, and ϕ_2 is the induced edge-map. The resulting graph morphism is clearly bijective.

As a trivial example of locally surjective graph morphism we have $\psi : \text{Star}_4 \rightarrow \text{Star}_3$, as shown in figure 3. Here ϕ_1 is defined by $\phi_1(0) = 0$, $\phi_1(1) = 1$, $\phi_1(2) = 2$, $\phi_1(3) = 2$.

Definition 3. Let $[F_Z, \sigma]$ and $[F_Y, \pi]$ be two SDS. An SDS-morphism between $[F_Z, \sigma]$ and $[F_Y, \pi]$ is a pair (ϕ, Φ) where $\phi : Y \rightarrow Z$ is a graph morphism and where $\Phi : \Gamma[F_Z, \sigma] \rightarrow \Gamma[F_Y, \pi]$ is digraph morphism.

FIGURE 3. The graphs Star_3 and Star_2 .

Given a graph morphism $\phi : Y \rightarrow Z$ we want to relate the dynamics of SDS over the two graphs Y and Z . The local functions will be the same for the two graphs unless otherwise stated. To begin, we relate update schedules for Y and Z via ϕ . Assume $|v[Y]| = n$ and $|v[Z]| = m$ and let $\phi^{-1}(i) = \{i_1, \dots, i_{l_i}\}$ where $i_1 < \dots < i_{l_i}$ for $1 \leq i \leq m$. Define the map $\eta_\phi : S_m \rightarrow S_n$ by

$$(2.1) \quad \eta_\phi(\pi = (\pi_1, \pi_2, \dots, \pi_m)) = (\pi_{1_1}, \dots, \pi_{1_{l_{\pi_1}}}, \dots, \pi_{m_1}, \dots, \pi_{1_{l_{\pi_m}}}).$$

For instance, in the example with $\phi : Q_2^3 \rightarrow K_4$, we have $\eta_\phi(4, 3, 2, 1) = (3, 4, 2, 5, 1, 6, 0, 7)$.

Similarly, we define the map $\tau : \mathbb{F}_2^m \rightarrow \mathbb{F}_2^n$ by

$$(2.2) \quad (\tau(x))_k = x_{\phi(k)}.$$

The dynamics of SDS over Y and Z can now be related in the following way [12]:

Theorem 1. *Let Y and Z be loop-free connected graphs, let $\phi : Y \rightarrow Z$ be a locally bijective graph morphism, and let $(f_i)_i$ be a fixed sequence of Boolean symmetric functions. Then the map τ induces a natural embedding*

$$(2.3) \quad T : \Gamma[F_Z, \pi] \hookrightarrow \Gamma[F_Y, \eta_\phi(\pi)].$$

Example 3. To illustrate the implications of Theorem 1 we show how to relate the phase space of $[\mathbf{Min}_{K_4}, \text{id}_4]$ to that of $[\mathbf{Min}_{Q_2^3}, (0, 7, 1, 6, 2, 5, 3, 4)]$. From the example above we have the bijective graph morphism $\phi : Q_2^3 \rightarrow K_4$. Next note that $\eta_\phi(\text{id}_4) = (0, 7, 1, 6, 2, 5, 3, 4)$. From, e.g., [1] we know that $[\mathbf{Min}_{K_4}, \text{id}_4]$ has exactly two 5-cycles and no fixed

points. The two 5-cycles are shown in the top row of figure 4. For convenience we use the map

$$\xi_i : \mathbb{F}_2^i \longrightarrow \mathbb{N}, \quad \xi_i(x_1, \dots, x_i) = \sum_{j=0}^i x_j \cdot 2^{j-1}$$

to encode states (binary tuples), and we have, e.g., $(1, 1, 0, 1) \mapsto 1 + 2 + 8 = 11$. It is straightforward to see that the phase space of $[\mathbf{Min}_{K_4}, \text{id}_4]$ is indeed embedded in the phase space of $[\mathbf{Min}_{Q_3^3}, (0, 7, 1, 6, 2, 5, 3, 4)]$.

We remark that $[\mathbf{Min}_{Q_2^3}, \eta_\phi(\text{id}_4)]$ has two fixed points in addition to the two 5-cycles shown in the last row in figure 4. These fixed points are related by the graph automorphism $\gamma = (07)(16)(25)(34)$, and consequently, so are their transients. Stated differently, the two components in $\Gamma[\mathbf{Min}_{Q_2^3}, \eta_\phi(\text{id}_4)]$ containing the fixed points are isomorphic. Their structure is shown in figure 5.

In view of example 3 it is of interest to determine the locally bijective graph morphisms $\psi : Y \rightarrow Z$ for a given graph Y . In the particular case of a locally bijective graph morphism $\psi : Q_2^n \rightarrow K_{n+1}$ we have that it exists if and only if a specific number theoretic condition is fulfilled.

Proposition 1. *Assume $2^n \equiv 0 \pmod{n+1}$, and let $\pi \in S_{n+1}$. Then there exists a covering $\phi : Q_2^n \rightarrow K_{n+1}$ and the SDS $[\mathbf{Par}_{Q_2^n}, \eta_\phi(\pi)]$ has a periodic orbit of length $n+2$.*

Proof. We will establish the existence of the covering map $\phi : Q_2^n \rightarrow K_{n+1}$ under the above condition in the next section. Since a covering map is locally bijective, we can apply Theorem 1 to deduce that the phase space of $\Phi = [\mathbf{Par}_{K_n}, \pi]$ can be embedded into the phase space of $\Psi = [\mathbf{Par}_{Q_2^n}, \eta_\phi(\pi)]$. Thus we see that *whatever we can deduce about the smaller system Φ applies to the larger system Ψ* . The phase space of Ψ has 2^{2^n} points while that of Φ has 2^{n+1} points.

One particular consequence of this is that every periodic orbit for Φ will also be a periodic orbit for Ψ . We will show that Φ always have a periodic orbit of length $n+2$. For simplicity we take $\pi = \text{id}_{n+1}$.




FIGURE 4. The top row shows the two five-cycles in $[\mathbf{Min}_{K_4}, \text{id}]$. The second row shows the images of the top cycles under χ_ϕ , and the last row shows the corresponding periodic cycles in the digraph $\Gamma[\mathbf{Min}_{Q_2^3}, \eta_\phi(\text{id}_4)]$.

By inspection $\text{Par}_n : \mathbb{F}_2^n \rightarrow \mathbb{F}_2^n$ is seen to satisfy the functional relation

$$(2.4) \quad \psi(x_1, \dots, x_{n-1}, \psi(x_1, \dots, x_n)) = x_n.$$




FIGURE 5. The structure of the components in $\Gamma[\mathbf{Min}_{Q_2^3}, \eta_\phi(\text{id}_4)]$ containing a fixed point. A single filled circle depicts a single state, while a circled number i depicts that there are i direct predecessors that do not have any successors themselves.

As a consequence of this we derive

$$\begin{aligned}
 x = (x_1, x_2, \dots, x_n) &\xrightarrow{1} (\text{Par}_n(x), x_2, x_3, \dots, x_n) \\
 &\xrightarrow{2} (\text{Par}_n(x), \text{Par}_n(\text{Par}_n(x), x_2, \dots, x_n), x_3, \dots, x_n) \\
 &= (\text{Par}_n(x), x_1, x_3, \dots, x_n) \\
 &\vdots \\
 &\xrightarrow{n} (\text{Par}_n(x), x_1, x_2, \dots, x_{n-1}),
 \end{aligned}$$

where \xrightarrow{i} denotes the update of state x_i .

In light of the above we obtain the commutative diagram

$$(2.5) \quad \begin{array}{ccc}
 \mathbb{F}_2^n & \xrightarrow{[\text{Par}_{K_n}, \text{id}]} & \mathbb{F}_2^n \\
 \downarrow \iota_{\text{Par}_n} & & \uparrow \text{proj} \\
 \hat{\mathbb{F}}_2^n & \xrightarrow{\sigma_{n+1}} & \hat{\mathbb{F}}_2^n,
 \end{array}$$

where

$$(2.6) \quad \text{proj}(x_1, \dots, x_n, x_{n+1}) = (x_1, \dots, x_n),$$

$$(2.7) \quad \iota_{\text{Par}_n}(x_1, \dots, x_n) = (x_1, \dots, x_n, \text{Par}_n(x_1, \dots, x_n)),$$

$$(2.8) \quad \sigma_{n+1}(x_1, x_2, \dots, x_{n+1}) = (x_{n+1}, x_1, \dots, x_n),$$

$$\text{and } \hat{\mathbb{F}}_2^n = \{x \in \mathbb{F}_2^{n+1} \mid x_{n+1} = \text{par}_n(x_1, \dots, x_n)\}.$$

Note that $\text{proj} : \hat{\mathbb{F}}_2^n \rightarrow \mathbb{F}_2^n$ and $\iota_{\text{Par}_n} : \mathbb{F}_2^n \rightarrow \hat{\mathbb{F}}_2^n$ are inverses. Similarly we obtain $[\text{Par}_{K_n}, \text{id}]^{(2)}(x) = (x_n, \text{Par}_n(x), x_1, x_2, \dots, x_{n-2})$ and in general $[\text{Par}_{K_n}, \text{id}]^{(k)} = \text{proj} \circ \sigma_{n+1}^k \circ \iota_{\text{Par}_n}$ showing that the order of an orbit of $[\text{Par}_{K_n}, \text{id}]$ is a divisor of $n + 1$.

Along the same lines we can show that $[\text{Par}_{K_n}, \pi]$ and $[\text{Par}_{K_n}, \text{id}]$ are topologically conjugated systems. We omit the details.

Thus we have that the length of every orbit of $[\text{Par}_{K_{n+1}}, \pi]$ is a divisor of $n + 2$. However, it is easy to see that the orbit containing the state $(1, 0, 0, \dots, 0)$ always have length $n + 2$. To be explicit we have for $n = 7$

$$\begin{array}{ccccccc} (1000000) & \longrightarrow & (1100000) & \longrightarrow & (0110000) & \longrightarrow & (0011000) \\ \uparrow & & & & & & \downarrow \\ (0000001) & \longleftarrow & (0000011) & \longleftarrow & (0000110) & \longleftarrow & (0001100) . \end{array}$$

Thus we deduce from Theorem 1 that $[\mathbf{Par}_{Q_2^n}, \eta_\phi(\pi)]$ has a periodic orbit of length $n + 2$, and we are done.

3. COVERING MAPS OVER THE n -CUBE

Proposition 1 turns out to be a special case of a more general theory for constructing covering maps over n -cubes. The theory for n -cubes, which may also extend to, e.g., certain Cayley graphs, is facilitated by the fact that \mathbb{F}_2^n can be viewed both as a group and as a vector space.

In this section we will show how the existence of subgroups $H < \mathbb{F}_2^n$ with certain properties can be used to obtain covering maps. In fact, this leads to a cascade of sequences of covering maps.

The following two results constitute the core of this theory[12]:

Lemma 1. *For any subgroup $H' < \mathbb{F}_2^n$ with $[\mathbb{F}_2^n : H'] \geq n + 1$ there exists an isomorphic subgroup $H \cong H'$ that has the property $H(x) \cap H(y) = \emptyset$ for $x \neq y$; $x, y \in \{0, e_1, \dots, e_n\}$.*

The following proposition shows how the subgroups of Lemma 1 induce covering maps in a natural way:

Proposition 2. *For each subgroup $H < \mathbb{F}_2^n$ with the property $H(x) \cap H(y) = \emptyset$ for $x \neq y$; $x, y \in \{0, e_1, \dots, e_n\}$ the graph $H \setminus \mathbb{Q}_2^n$ is connected, undirected and loop-free and the natural projection*

$$\pi_H : \mathbb{Q}_2^n \longrightarrow H \setminus \mathbb{Q}_2^n, \quad v \mapsto H(i)$$

is a covering map.

Proof. We have to show that the π_H -induced restriction mapping:

$$(3.1) \quad \text{res}_{\text{Star}_{\mathbb{Q}_2^n}(\xi)}(\pi_H) : \text{Star}_{\mathbb{Q}_2^n}(\xi) \longrightarrow \text{Star}_{H \setminus \mathbb{Q}_2^n}(\pi_H(\xi))$$

is an isomorphism for arbitrary $\xi \in \mathbb{F}_2^n$. By construction, $\text{res}_{\text{Star}_{\mathbb{Q}_2^n}(\xi)}(\pi_H)$ is surjective and $H(x) \cap H(y) = \emptyset$ for $x \neq y$; $x, y \in \{0, e_1, \dots, e_n\}$ is equivalent to $H(x + \xi) \cap H(y + \xi) = \emptyset$ for $x \neq y$; $x, y \in \{0, e_1, \dots, e_n\}$ for any $\xi \in \mathbb{F}_2^n$. Therefore $\text{res}_{\text{Star}_{\mathbb{Q}_2^n}(\xi)}(\pi_H)$ is injective and the proof of the proposition is complete.

Corollary 1. *Let n be a natural number. Then we have $2^n \equiv 0 \pmod{n+1}$ if and only if there exists a subgroup $G < \mathbb{F}_2^n$ with the property $\mathbb{F}_2^n = G(0) \cup \bigcup_{i=1}^n G(e_i)$.*

Proof. Suppose we have $2^n \equiv 0 \pmod{n+1}$. Since \mathbb{F}_2^n is a p -group there exists a subgroup in its decomposition series $H < \mathbb{F}_2^n$ with the property $[\mathbb{F}_2^n : H] = n + 1$. According to Lemma 1, there exists some set of H -representatives $\{\varphi_1, \dots, \varphi_n\}$ that forms a basis of \mathbb{F}_2^n . Let f be the \mathbb{F}_2 -homomorphism defined by $f(\varphi_i) = e_i$, for $i = 1, \dots, n$. Clearly, $G = f(H)$ has the property $\mathbb{F}_2^n = G(0) \cup \bigcup_{i=1}^n G(e_i)$, whence the corollary.

Corollary 2. *There exists a locally bijective graph morphism*

$$(3.2) \quad \varphi : \mathbb{Q}_2^n \longrightarrow K_{n+1}$$

if and only if $2^n \equiv 0 \pmod{n+1}$ holds.

The following lemma establishes a group action of $\text{Aut}_0(Q_2^n) = \{\sigma \in \text{Aut}(Q_2^n) \mid \sigma(0) = 0\}$ on graphs $H \setminus Q_2^n$.

Lemma 2. *Let $H < \mathbb{F}_2^n$ be a subgroup and let $\eta \in \text{Aut}_0(Q_2^n)$. Then we have*

$$H \setminus Q_2^n \cong \eta H \setminus Q_2^n .$$

3.1. Computation of covering maps. For small n we can easily compute the various subgroups of $Q_2^n[4]$. It turns out that for fixed $n \leq 7$ the orbits of the subgroups under $\text{Aut}_0(Q_2^n)$ (the isometric orbits) yield non-isomorphic reduced graphs. Moreover, for $n = 4$ the only covering map images of Q_2^n are the ones that can be obtained as $H \setminus Q_2^n$. One may speculate if this is true in general. That is, do the isometric orbits yield non-isomorphic covering images and can all covering images be obtained as $H \setminus Q_2^n$ for some sub-group H .

In table 1 we have summarized the number of non-isometric orbits for all subgroups of Q_2^n satisfying the condition in Proposition 2 for $n \leq 9$. In table 2 orbit representatives are given for $n \leq 7$. Some of the reduced graphs occurring in this scheme are shown in figure 6 and 7.




FIGURE 6. The only two non-isomorphic graphs of the form $H \setminus Q_2^4$.

3.2. Constructing covering graphs from chain maps. In [6] a way of constructing graph coverings of a given graph is described. We




FIGURE 7. All non-isomorphic graphs of the form $H \setminus Q_2^7$ on 16 vertices. In each case there are two interlocked cubes. The edges in red show how the two cubes are connected. The other vertices of the outer cube are similarly connected, but the edges are not shown to ease visualization. Note that there are 4 lines in figure on the top right ★. The two lines connecting the outer cube to the closest vertex and the most distant vertex of the inner cube coincide.

will outline it briefly. Let $S\Gamma$ denote the set of arcs or sides of the graph Γ . Thus each edge $\{u, v\}$ gives rise to two sides, (u, v) and (v, u) . Let G be any group. A G -chain on Γ is a map $\phi : S\Gamma \rightarrow G$

n	#(size 2)	#(size 4)	#(size 8)	#(size 16)
4	2	0	0	0
5	3	1	0	0
6	4	4	1	0
7	5	8	5	1
8	6	14	15	6
9	7	22	38	-

TABLE 1. Subgroups of the n -cube: The table gives the number of isomorphic-non-isometric orbits of subgroups $H < Q_2^n$ of the given sizes.

such that $\phi(u, v) = (\phi(v, u))^{-1}$ for all sides (u, v) of Γ . The *covering map* $\hat{\Gamma} = \hat{\Gamma}(G, \phi)$ of Γ with respect to a given G -chain ϕ on Γ is the graph with $v[\hat{\Gamma}] = G \times v[\Gamma]$ and where vertices (g_1, v_1) and (g_2, v_2) are joined by an edge iff $(v_1, v_2) \in S\Gamma$ and $g_2 = g_1\phi(v_1, v_2)$. $\hat{\Gamma}$ is clearly well-defined.

Note that the 3-cube is isomorphic to the covering graph $\hat{K}_4(\mathbb{F}_2, \phi)$ of K_4 where ϕ is the \mathbb{F}_2 -chain assigning 1 to each side of K_4 .

Moreover we note that the graph G_5 on the third row in figure 7 is isomorphic to the covering graph $\hat{K}_8(\mathbb{F}_2, \phi)$, where ϕ again is the \mathbb{F}_2 -chain assigning 1 to each side of K_8 .

Another, but quite similar, approach for constructing covering maps uses so-called “voltage graph” [7, 8, 9]. Voltage graphs and G -chains are strongly related.

4. SYNOPSIS

In this paper we have introduced a mathematical framework suitable for the analysis of network dynamics. A new class of dynamical systems, SDS, that is an abstract model for the dynamics of networks has been discussed.

The results on SDS presented in this paper display the interplay between the structure of the underlying networks and their corresponding

n	Size 2	Size 4	Size 8	Size 16
4	{0, 14}, {0, 15}			
5	{0, 28}, {0, 30}, {0, 31}	{0, 7, 25, 30}		
6	{0, 56}, {0, 60}, {0, 62}, {0, 63}	{0, 7, 25, 30}, {0, 7, 56, 63}, {0, 7, 57, 62}, {0, 15, 51, 60}	{0, 7, 25, 30, 42, 45, 51, 52}	
7	{0, 112}, {0, 120}, {0, 124}, {0, 126}, {0, 127}	{0, 7, 25, 30}, {0, 7, 56, 63}, {0, 7, 57, 62}, {0, 7, 120, 127}, {0, 7, 121, 126}, {0, 15, 51, 60}, {0, 15, 113, 126}, {0, 15, 115, 124}	{0, 7, 25, 30, 42, 45, 51, 52}, {0, 7, 25, 30, 97, 102, 120, 127}, {0, 7, 25, 30, 98, 101, 123, 124}, {0, 7, 25, 30, 106, 109, 115, 116}, {0, 15, 51, 60, 85, 90, 102, 105}	{0, 7, 25, 30, 42, 45, 51, 52, 75, 76, 82, 85, 97, 102, 120, 127}

TABLE 2. Subgroups of the n -cube: The table gives representatives of the isomorphic-non-isometric orbits of subgroups $H < Q_2^n$ of the given sizes that satisfy the condition in Lemma 1. Explicit computations show that the orbits give non-isomorphic reduced graphs $H \setminus Q_2^n$ for $n \leq 7$.

dynamics. Theorem 1 allows one, for particular schedules, to obtain key information about a given system via a reduced/simpler network. We have shown for the special case of Boolean n -cubes how to construct these reduces systems based on the subgroup structure of \mathbb{F}_2^n . Networks over n -cubes can be reduced to networks over smaller graphs induced by symmetries of the n -cube itself.

ACKNOWLEDGEMENTS

We thank C.L. Barrett for stimulating discussions. Special thanks and gratitude to D. Morgeson for his continuous support. This research is supported by Laboratory Directed Research and Development under DOE contract W-7405-ENG-36 to the University of California for the operation of the Los Alamos National Laboratory.

REFERENCES

- [1] C. L. Barrett, H. S. Mortveit, and C. M. Reidys. Elements of a theory of simulation II: Sequential dynamical systems. *Appl. Math. and Comput.*, 107(2-3):121–136, 2000.
- [2] C. L. Barrett, H. S. Mortveit, and C. M. Reidys. Elements of a theory of simulation III: Equivalence of SDS. *Appl. Math. and Comput.*, 2000. In press (2000).
- [3] C. L. Barrett, H. S. Mortveit, and C. M. Reidys. Elements of a theory of simulation IV: Fixed points, invertibility and equivalence. *Appl. Math. and Comput.*, 2000. Submitted (2000).
- [4] C. L. Barrett, H. S. Mortveit, and C. M. Reidys. Elements of a theory of simulation V: Phase space properties of sequential dynamical systems. *Appl. Math. and Comput.*, 2001. In progress (2001).
- [5] C. L. Barrett and C. M. Reidys. Elements of a theory of simulation I: Sequential CA over random graphs. *Appl. Math. and Comput.*, 98:241–259, 1999.
- [6] N. Biggs. *Algebraic Graph Theory*. Cambridge Mathematical Library, second edition, 1993.
- [7] J. L. Gross. Voltage graphs. *Discrete Mathematics*, 9:239–246, 1974.
- [8] J. L. Gross and T. W. Tucker. Generating all graph coverings by permutation voltage assignments. *Discrete Mathematics*, 18:273–283, 1977.
- [9] M. Hofmeister. Isomorphisms and automorphisms of graph coverings. *Discrete Mathematics*, 98:175–183, 1988.
- [10] H. S. Mortveit and C. M. Reidys. Discrete, sequential dynamical systems. *Discrete Mathematics*, 226:281–295, 2001.
- [11] C. M. Reidys. On Acyclic Orientations and SDS. *Adv. in Appl. Math.*, 2000. Accepted (2000).
- [12] C. M. Reidys. Phase space properties of SDS. *Adv. in Appl. Math.*, 2001. In progress (2001).

LOS ALAMOS NATIONAL LABORATORY, D-2, 87545 NEW MEXICO, USA, MAIL-
STOP: TA-0, SM-1237, MS M997, PHONE: ** (505) 665-7285, FAX: ** (505)
665-7464

E-mail address: `henning@lanl.gov`

LOS ALAMOS NATIONAL LABORATORY, TSA-2, 87545 NEW MEXICO, USA,
MAILSTOP: TA-0, SM-1237, MS M997, PHONE: ** (505) 665-0911, FAX:
** (505) 665-7464

E-mail address: `duck@tsasa.lanl.gov`